

HEADQUARTERS  
JOINT TASK FORCE SEVEN  
APO 187 (HOW) c/o POSTMASTER  
SAN FRANCISCO, CALIFORNIA

A57

J-3/7293

18 March 1954

SUBJECT: Radiological Surveys of Several Marshall Island Atolls

TO: Distribution

OK UmeL  
RDK  
OK UmeL  
PL  
3-9-54

1. Attached herewith for your information and retention are copies of radiological surveys made on certain Marshall Island Atolls. The surveys were conducted as a result of contamination deposited on the affected atolls by BRAVO Shot, Operation CASTLE, fired from a reef approximately one and one half nautical miles southwest of Namu, Bikini Atoll. BRAVO Shot time was 1845 Zebra, 28 February 1954.

2. Water and soil samples were shipped to the Health and Safety Laboratory, New York Operations Office, Atomic Energy Commission (Attention: Mr. Merrill Eisenbud) for analysis.

FOR THE COMMANDER:

DISTRIBUTION:

- CTG 7.1 - Copy 1-30
- CTG 7.2 - Copy 31
- CTG 7.3 - Copy 32
- CTG 7.4 - Copy 33
- CTG 7.5 - Copy 34
- CINCPAC - Copy 35
- CINCPACFLT - Copy 36
- HICOMTERPACIS - Copy 37
- COMNAVSTAKWAJ - Copy 38
- DM/REC - Copy 39
- DE/REC - Copy 40
- Ch AFSWP - Copy 41
- CG FldComd(DWET) - Copy 42
- C/S USA, ExAgt - Copy 43
- LASL H Div. - Copy 44
- HASL, NYOO (c/o Mgr Opns) - Copy 45-46
- USS RENSHAW (DDE-499) - Copy 47
- USS PHILIP (DDE-498) - Copy 48
- USS NICHOLIS (DDE-449) - Copy 49

*E. McGinley*

E. MCGINLEY  
Brigadier General, U.S. Army  
Chief of Staff 326 US ATOMIC ENERGY COMMISSION

Location LANL  
Collection Records Center  
Folder CASTLE FALLOUT OBSERVATIONS

CONFIRMED TO BE UNCLASSIFIED

BY AUTHORITY OF DOE/OD

REVIEWED BY J. Mag 9/14/83  
DATE

3 Incls:

1. Report on Soil and Water Sampling Mission by Major R. D. Crea
2. Report on Soil and Water Sampling Mission by Dr. T. N. White, LASL
3. Rad. Survey of Downwind Atolls Contaminated by BRAVO by Dr. Herbert Schmitt

COPIED/DOE  
LANL RC

SRD-213-54E

~~NO~~  
~~44~~  
~~45~~  
~~46~~  
~~47~~  
~~48~~  
~~49~~

=====

15

101

HEADQUARTERS  
JOINT TASK FORCE SEVEN  
APO 187 (HOW) c/o POSTMASTER  
SAN FRANCISCO, CALIFORNIA

COMPT

8 March 1954

SUBJECT: Report on Soil and Water Sampling Mission

TO: Commander  
Joint Task Force SEVEN  
APO 187 (HOW)  
c/o Postmaster  
San Francisco, California

*OK over RDK*  
*OK under PL 3-9-54*

1. In compliance with your oral instructions, the undersigned visited LIKI and AILUK Atolls, JEMO Island and MEJIT Island in the Eastern Marshalls between the period 5-8 March 1954 for the purpose of collecting soil and water samples and measuring level of gamma radiation present at those places in connection with BRAVO. The mission, consisting of the undersigned and a Marshallese interpreter, Lan Lakapun, embarked on the USS RENSHAW (DDE499) at Kwajalein, visited the four sites and returned to Bikini, where the remainder of the trip to Eniwetok was performed by PBM. There follows a detailed discussion of the findings at each location:

a. LIKIEP ATOLL. The samples were taken on Likiep Island, which had the largest native population. Access to the lagoon was gained through South Pass. Poor light at the end of the day and numerous coral heads necessitated anchoring about 4 miles from Likiep Island. Trip in was made by whaleboat the following morning. A water sample was taken from a large cistern fed from the roof of the Catholic rectory, and earth samples were taken from random spots about the island which were unsheltered by trees or other growth at approximately 0800 M 6, March 1954. Radiation readings were taken with a MX-5 instrument between 0800 M and 0900 M and showed a maximum of 3 milliroentgens per hour. No variations from this reading were noted on clothing or bare feet of individuals. According to accounts received by Bishop Feeney, S.J., the population was greatly excited by the light and blast wave, the latter which reportedly arrived about 30 minutes subsequent to the light flare. According to Bishop Feeney, church attendance was greatly stimulated on the day of the test.

b. JEMO Island. This location was reached at 1100 M, 6 March 1954. It consists of a small heavily wooded island, surrounded by a line coral reef with heavy surf on three sides. There being no place for landing a whaleboat, personnel and equipment were transferred from the whaleboat to the reef by a one man rubber raft. The undersigned transferred himself by swimming. The island proved to be uninhabited, and reportedly is a sea turtle preserve. Turtle hunters erected several houses, a rain barrel of which provided a water sample. Earth samples were gathered at random from open areas, including one of beach sand above the high tide mark. The party was led straight across the island and back to the landing area via the beach, in order to verify its uninhabited state. Samples were

COPIED/DOE  
LAW/RC

44-85-2 (PAGES)  
B

RESTRICTED USE ONLY

10-10-54  
16  
497

CONFIDENTIAL

SUBJECT: Report on Soil and Water Sampling Mission

collected at approximately 1200 M, 6 March 1954. Instrument readings with the MX-5 showed a maximum of 3 mr/hr, however this was not considered reliable, since a higher scale showed a lower reading.

c. AILUK ATOLL. The ship reached this atoll at approximately 1600 M, 6 March 1954, and slowly moved to an anchorage off Ailuk Island, the most heavily populated. The lagoon has not been swept, and numerous coral heads and pinnacles provided considerable hazard to ship movement. The landing party moved ashore by whaleboat without difficulty, and again obtained water samples from the most prominent cistern and soil samples from random unsheltered spots. Readings with the MX-5 showed approximately 3 mr/hr (off the 2 mr scale). An AN/PDR-27E showed a high reading of 7 mr/hr, however, on a different scale a reading of 12 or 15 mr/hr was obtained. The MX-5 reading is probably nearest correct. No significant variations were detected on bare feet or clothing of individuals. Samples and readings were taken at approximately 1700 M, 6 March 1954.

d. MEJIT Island. This single coral island is also surrounded by a reef, as is JEMO, but landing was possible with a whaleboat, due to an area protected from the surf. The island was found to be heavily populated in view of its size, the total number of people being 327, according to the island magistrate. Soil and water samples were taken as in the previously described manner, at approximately 1300 M, 7 March 1954. Readings with the MX-5 showed maximum of approximately 3 mr/hr (off the 2 scale, but approximately 1.5 on the 20 scale); the maximum reading with a PDR 27 E was 10 mr/hr. The true figure was probably somewhere between the two.

2. CONCLUSIONS. Low level (less than 10 mr/hr) radiation measurements with field instruments of the type used are highly unsatisfactory. One MX-5 and three AN/PDR 27 E instruments all showed widely variant readings on different scales, and varied among each other when exposed to the same radiation. An AN/PDR T1-B proved completely useless not holding to zero even after an hours warm-up, and also showing widely variant readings on different scales.

3. RECOMMENDATIONS. Landing parties in islands such as JEMO and MEJIT should be provided with a rubber 6-man or 8-man pneumatic boat, to provide greater safety to personnel and equipment. This will permit landing directly on live coral reefs with less danger of the boat being stove in. Ships assigned to such missions should draw such equipment prior to departure.

4. The successful accomplishment of the mission was greatly facilitated by the interest and enthusiasm of the Commanding Officer of the USS RENSHAW, CDR L. H. Alford, USN, and his officers and men. Their material contributions were necessary to the mission, however, the many valuable suggestions and assistance in solutions of problems proved invaluable.

COPIED/DOE  
LANL RC

/s/ R. D. Crea  
R. D. CREAA  
MAJ, USA

CONFIDENTIAL

MEMORANDUM FOR: CJTF SEVEN

10 March 1954

*on file  
3-9-53*

SUBJECT: Report on Soil and Water Sampling Mission

1. In compliance with your oral instructions, the undersigned visited Wotje, Erikub, Maloolap, Wotho, Majuro Atolls in the Marshall Islands 5 through 7 March 1954 for the purpose of obtaining earth and drinking water samples, and of measuring gamma ray dose rates, and also checked the radiological condition of the S.S. ROQUE on its arrival at Majuro 7 March 1954.

2. The first four atolls were visited by Marshallese interpreter Takushi and the writer by means of an UF-1 amphibious aircraft. Majuro was reached by C-47. Erikub might have been omitted since it was not inhabited, being property of the Wotje tribe which goes there only occasionally to gather copra. (This was unknown until after the visit.)

3. At each atoll, only the principal inhabited island was visited. At each visited island an effort was made to compose a representative soil average by collecting into a single container several samples, each approximately one square foot of area and one inch depth. Water samples were collected from the principal sources currently in use. The gamma dose rates are averages for the inhabited areas.

4. With regard to certain minor discrepancies between the survey methods used by Major R. D. Crea and the writer; it was originally planned to perform the survey jointly, and when it became advisable to separate and survey different atolls, no time remained for discussion of details of techniques.


5. Gamma-ray dose rates on Wotje and on Erikub are each the average of MX-5 and AN/PDR-39 average readings which agreed reasonably well. The MX-5 was rendered inoperative when the rubber life raft was swamped by surf on the first attempt to launch from the beach at Erikub. Following the Wotho survey, the PDR-39 developed a temperature-dependent reading of 0.4 - 2 mr/hr, so that later readings in this range are of very dubious reliability.

6. The following tabulation summarizes the atoll survey. S is Soil, W is Water Sample:

<u>ATOLL</u>	<u>ISLAND</u>	<u>DATE</u>	<u>TIME</u>	<u>SAMPLE NO</u>	<u>MR/HR &amp; SAMPLING</u>
WOTJE	ORMED	5 Mar	1600	S5	3.5 mr/hr, 1-beach, 3-mid-vill- age, 1-back village.
				W6	1/2 well plus 1/2 catch basin.
ERIKUB	ERIKUB	5 Mar	1715	S6	1.5 mr/hr. 1-mid-village, 1 on path to beach. No inhabit- ants, no water supply found.

COPIED/DOE  
LANL RC

THIS DOCUMENT CONTAINS (3) PAGES  
44 85 B



18

**RESTRICTED**

<u>ATOLL</u>	<u>ISLAND</u>	<u>DATE</u>	<u>TIME</u>	<u>SAMPLE NO</u>	<u>MR/HR &amp; SAMPLING</u>
MALOELAP	KAVEN	6 Mar	1130	S7	1.8 mr/hr, 2-villago, 2-path to beach.
				W12	Well water.
				W13	From catch basin.
WOTHO	WOTHO	6 Mar	1615	S8	0.8 mr/hr, 1 by well; 2-mid-villago.
				W9	Well water (no rain in catch basin for 2 mo.)
MAJURO	ULIGA	7 Mar	1200	S9	0.5 mr/hr, 4 from near Admin Bldg.
				W10	Tap water.

7. Pacific Micronesian Line S.S. "ROQUE", Master: Lawrence Blanc, home port, Guam, left Eboye 0840 M on 1 March, entered channel to Utirik Lagoon about 1200 M on 2 March, and anchored in Lagoon at 1524 M on 2 March; docked at Majuro (Uluga Is.) 1630M on 7 March. Readings (mr/hr) after docking: 2-3 inside main dock structure, 10 on open deck, 5-8 in sleeping quarters on upper deck, 10-30 on rope and canvas. Prior radiation levels cannot be estimated because of rain squalls and uncertainty about when decks last washed. Master was advised to have decks washed down as soon as convenient. He was told that the activity would not hurt anyone, but that it was undesirable to have it around longer than necessary.

8. RECOMMENDATIONS: Future visits to Erikub and Maloelap should not be attempted by UF-1 except under conditions of greater urgency. The writer's prior experience in such operations is very limited, but from his own observations plus the remarks made by those better qualified to judge, it appears that a fair amount of risk is involved.

9. Especially notable was the very cooperative attitude of the Navy personnel at Kwajalein and the Marshall District Administrative Officials at Majuro in supporting this mission.

1 Incl:  
 Marshall Islands Atoll  
 Samples collected by T. N.  
 White, 5-7 March 1954

/s/ T. N. White  
 DR. T. N. WHITE  
 Health Division  
 LASL

COPIED/DOE  
 LANL RC

=====

**RESTRICTED**

**RESTRICTED**

[REDACTED]

[REDACTED]

MARSHALL ISLANDS ATOLL SAMPLES COLLECTED BY T. N. WHITE, 5-7 MARCH 1954

Earth samples were collected as follows:

At each island visited several samples were dug and put into the same one-gallon "ice-cream carton". Each sample (i.e. each digging) approximated one square foot to a depth of one inch. The number and locations of the samples were selected to represent, as well as could be judged, an average of the areas used by the inhabitants, after the samples were mixed in the carton. Areas that were unusually shaded or unshaded by trees were avoided. The large "pebbles" in the composite represent coral gravel from "main street" through the village.

Water samples were selected according to the principal source in current use.

COPIED/DOE  
LANL PC

Inclosure 1

[REDACTED]

[REDACTED] - 1346

20

HEADQUARTERS TASK UNIT 13  
Task Group 7.1  
APO 1B7 (HOW) P.O. Box 8  
c/o Postmaster  
San Francisco, California

~~CONFIDENTIAL~~

OK  
PL  
3-9-53

TU-13-54-375

12 March 1954

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

1. Acknowledgment

The members of the survey team wish to express their appreciation to the Captain, officers and members of the crew of the USS NICHOLAS (DDE 449) for their assistance and cooperation in conducting the survey herein reported. Captain Elliot turned over all possible facilities of his ship in order to assist in the survey. LT Frink, the Executive Officer, organized all the operations of the boat parties, and it was only through his personal direction and participation that it was possible to carry out the small boat surveys under extremely difficult conditions. Since most of the lagoon waters were not navigable by a DDE, it was necessary to make long boat trips in high seas and land on tricky coral reefs. That it was possible to make, without mishap, a detailed survey of five widely separated atolls in the course of three days with only two boats was largely due to his efforts.

2. Introduction

The BRAVO Shot contaminated a number of atolls in generally eastward direction from Bikini to such an extent that it became necessary to evacuate the native populations from Rongelap, Ailinginae and Utirik Atolls and the military personnel on Rongerik Atoll. Following this evacuation CJTF SEVEN organized the subject detailed radiological survey of the atolls to the eastward of Bikini (Ref. CJTF SEVEN Eniwetok 060400Z). The data from this survey were required for the following purposes:

- a. The evaluation of the radiation effects on evacuees.
- b. The estimation of the elapsed time before reoccupancy.
- c. The estimation of the residual radiation effects of large yields surface detonations.

In connection with this survey, teams from various Task Groups and Mr. Wilds, Trust Territory Representative, returned to the atolls to secure the evacuated habitations, service military equipment, and obtain documentary photography.

3. Operational Schedule

8 March - 0800 Survey team rendezvous aboard USS NICHOLAS (DDE 449) in Rongelap Lagoon.

COPIED/DOE  
LANL RC

44 85 11 PAGE(S)  
B

~~CONFIDENTIAL~~

CR-912-Ed-2E

~~CONFIDENTIAL~~

21  
173

~~CONFIDENTIAL~~

~~CONFIDENTIAL~~

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

- 8 March - 1000 - 1800 Two parties in small boats surveyed living areas on Rongelap Island and eastern half of Rongelap Atoll.
- 9 March - 0700 - 1130 Two parties in small boats proceeded from the DDE which was stationed outside Utirik Atoll and surveyed Utirik and Aon Islands, the main islands of the Atoll.
- 9 March - 1500 - 1700 One party in a small boat landed on the outer reef of Bikar Island and surveyed the island, the only large island of Bikar Atoll.
- 10 March - 0700 - 1100 Two parties in small boats proceeded from the DDE which was stationed outside Rongerik Atoll and surveyed Eniwetak Island (where the Task Force's Units had been stationed) and the other important islands of the Atoll.
- 10 March - 1430 - 1900 Two parties in small boats proceeded from the DDE which was stationed outside Alinginae Atoll and surveyed the inhabited islands of the Atoll.
- 11 March - 0700 - 1400 One party in a small boat surveyed the northwestern islands of Rongelap Atoll and one party rechecked the living areas on Rongelap Island and established a reference location for future decay measurements.
- 12 March - 0800 Survey team arrived Eniwetok Atoll via DDE.

4. The following personnel from test projects in TG 7.1, TU 13, served as members of the survey team:

Herbert Scoville, Jr.	TU-13 Staff
Richard Rast	Project 2.1
Richard Soule	Project 2.5a
Walmer Strobe	Project 6.4

The USS NICHOLAS (DDE 449) supplied boat crews under the direction of LT Clifford Frink, Executive Officer, for surveys.

5. Instrumentation

Radiac set AN/PDR-39 was selected as the instrument to be used in the conduct of the survey. Five (5) each of AN/PDR-39 were calibrated with an 80 Curie Co<sup>60</sup> source twenty-four hours before departure. The calibration yielded a zero variation between instruments - any scale. Upon cross checking three of these instruments, (a point of actual survey) in a radiation field of 0.320 r/hr it was found that all three instruments gave the same reading.

COPIED/DOE  
LANL RC ✓

These survey meters were subject to prolonged use under adverse conditions of dampness (to the point of sea water splashing over them), salt deposit and continual rough handling. With one exception, all instruments operated efficiently for the duration of the operation. On the final day it was found

~~CONFIDENTIAL~~

~~CONFIDENTIAL~~

27

~~CONFIDENTIAL~~

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

that one survey meter could not be properly zero adjusted. The four remaining AN/PDR-39, still operated efficiently and seemed to be in good working order.

One (1) each Bockman MX-5, and one (1) each AN/PDR-27A was brought along for any low intensity checks necessary. Two (2) each calibrated AN/PDR-TLB, were on hand to serve as spares in the event of operational failure with the AN/PDR-39. None of these instruments were required.

6. The average and maximum gamma dose rates measured on the various islands of each atoll are plotted in Figures 1 through 5. All measurements were made at waist height unless otherwise indicated. The maximum readings do not include measurements made with the instrument next to a contaminated surface.

Detailed surveys were made of all the inhabited localities. Typical readings are given in Tables 1 and 2 for the native village of Rongelap Island, and the TG 7.4 camp on Eniwetak Island. In general, the villages and the camps appeared to have slightly lower average dose rates than the remainder of the island. This can perhaps be ascribed to different geometry of the contamination and to slightly greater penetration into the loose gravel in the native villages. The dose rates inside the native huts appeared to be almost the same as the dose rate outside. The dose rate in the middle of the military barracks, tents, and shacks was 1/3 to 1/2 that outside. This reduction is probably largely a geometrical effect. The dose rate fell off rapidly on the beach below the high tide mark. There was no evidence of rain washing off the contaminated material. The foliage on the windward sides of the islands appeared to be slightly above average contamination.

TABLE 1

TYPICAL READINGS IN RONGELAP VILLAGE - 8 MARCH

<u>Location</u>	<u>Dose Rate (mr/hr)</u>
Rongelap Island (average)	375
Center of village	280
Near central cistern	300
Near southern cistern	220
Near northern cistern	350

TABLE 2

TYPICAL READINGS IN CAMP ON ENIWETAK IS. - 10 MARCH

<u>Location</u>	<u>Outside Dose Rate (mr/hr)</u>	<u>Inside Dose Rate (mr/hr)</u>
Eniwetak Island (average)	280	--
Mess hall	220	110
Tent, edge of main camp	270	175
Latrine	260	160
Sleeping quarters	260	90
Dispensary	220	110

COPIED/DOE  
LANL RC

23

~~CONFIDENTIAL~~

=====

**RESTRICTED - USE ONLY**

Radio Station	290	160
Weather Station (N end of island)	280	110
Proj 6.6. Station (S end of island)	240	--

In order to estimate the rate of decay between 8 and 11 March, the following radiation measurements were taken on three days on Rongelap Island:

	<u>8 March</u>	<u>11 March</u>
Central living area (village)	280 mr/hr	170 mr/hr
Southern most cistern	220 mr/hr	145 mr/hr
Roof of cistern (Southern most)	240 mr/hr	140 mr/hr
Ground (contact) cistern area	220 mr/hr	110 mr/hr

An area was selected 30 yards inland from the Rongelap cemetery as a measuring point for future decay measurements. This area is outlined with 2X4s placed on pails. The waist height reading was 210 mr/hr at 1000 hours, 11 March 1954.

### 7. Sample collections

Water samples were collected from the water supplies of all inhabited areas. About two quarts of water were transferred to a polyethylene bottle at each site. These will be turned over to the New York Operations Office, AEC for analysis.

Soil samples were collected at all inhabited areas and also at several uninhabited islands. In collecting the soil samples a one foot by one foot square was marked on the ground and soil to about one inch of depth was removed from the square and transferred to a cardboard container. The primary samples will be turned over to the New York Operation Office, AEC, for analysis, and some smaller samples will be analyzed by Program 2 of TU 13.

Listed in Table 3 are the samples taken with the dose rate measured at waist height at the location where they were taken.

TABLE 3 - SOIL

<u>Sample No.</u>	<u>Atoll</u>	<u>Island</u>	<u>Date</u>	<u>Mr/Hr</u>
1*	Rongelap	Rongelap (North end)	8 Mar	440
2	Rongelap	Rongelap (Center of village)	8 Mar	280
3	Rongelap	Rongelap (1 mile north of village)	8 Mar	340
4	Rongelap	Rongelap (near South cistern of village)	8 Mar	220
5*	Rongelap	Eriirippu	8 Mar	2200
6*	Rongelap	Eniaetok	8 Mar	900
7*	Rongelap	Kabelle	8 Mar	2000
8*	Utirik	Utirik	9 Mar	40
9	Bikar	Bikar	9 Mar	160
10	Rongerik	Ehiwotak	10 Mar	280
11*	Ailinginae	Sifo	10 Mar	100

COPIED/DOE  
LANL RC

\*Small additional sample taken for analysis by Program 2 of TU 13.

**RESTRICTED - USE ONLY**

24

**OFFICIAL USE ONLY**

TABLE 9 - WATER

<u>Sample No.</u>	<u>Atoll</u>	<u>Island</u>	<u>Date</u>	<u>Mr/Hr</u>
1	Rongelap	Rongelap (central cistern)	8 Mar	300
2	Rongelap	Rongelap (North part of village)	8 Mar	350
3	Rongelap	Rongelap (Northernmost cistern)	8 Mar	400
4	Rongelap	Rongelap (Southernmost cistern)	8 Mar	220
5	Utirik	Utirik (cistern near church)	9 Mar	40
6	Utirik	Utirik (cistern at south of village)	9 Mar	40
7	Rongerik	Eniwetak (Distillation water)	10 Mar	240

In addition to the above, a sample of foliage was taken at the windward side of Bikar Island. The radiation field was 180 mr/hr on 9 March 1954 at this point.

8. Conclusions and Recommendations

a. The radiological survey proved that a large yield surface detonation can produce extremely serious radiological contamination over a distance more than 120 miles downwind and important contamination about 250 miles downwind.

b. The center of the contamination pattern from the BRAVO Shot lies somewhat north of Rongelap and Rongerik Atolls and probably not far from a line between Bikini and Bikar.

c. Although the fall-out was serious on Rongelap Island located at the extreme southeast tip of the atoll, the contamination was about ten times greater at the north side of the atoll, twenty miles away.

d. The contamination decreased by a factor of about eight over the downwind distance of 50 miles between Rongelap and Rongerik.

e. Standard military field housing provides a significant degree of protection to personnel inside.

f. The AN/PDR-39 proved to be a very satisfactory instrument for field survey work under rigorous environmental conditions.

g. A single DDE with two (2) whale boats is not a completely satisfactory method of conducting a broad radiological survey of the type just completed. Future surveys should consider using vessels capable of entering more of the atolls and of handling a helicopter and several small boats.

COPIED/DOE  
LANL RC

6 Incls:

- 1. Rad. Survey Rongelap
- 2. Rad. Survey Utirik
- 3. Rad. Survey Bikar
- 4. Rad. Survey Rongerik
- 5. Rad. Survey Ailinginae
- 6. Summary of Rad. Survey

/s/ Herbert Scoville  
DR. HERBERT SCOVILLE  
Technical Director  
AFSWP

**RESTRICTED**


~~RESTRICTED~~





~~SECRET~~

~~SECRET~~  
~~CONFIDENTIAL~~  
~~CONFIDENTIAL~~



**BIKAR ATOLL  
RADIOLOGICAL SURVEY**

9 MARCH  
AVE. READING (MAX. READING)  
MR / HR

~~SECRET~~  
~~CONFIDENTIAL~~  
~~CONFIDENTIAL~~


**FIG. 3**

COPIED/DOE  
LANL RC

~~SECRET~~

2.1

[REDACTED]



RONGERIK ATOLL  
RADIOLOGICAL SURVEY  
10 MARCH  
AVE READING (MAX. READING)  
MR / HR

FIG. 4

COPIED/DOE  
LANL RC ✓

[REDACTED]

AILINGINAE ATOLL  
 RADIOLOGICAL SURVEY

10 MARCH

AVERAGE READING (MAX. READING)  
 MR/HR.




FIG. 5

SUMMARY OF RADIOLOGICAL SURVEY

NOTE

ALL VALUES ARE R/HR AT REFERENCE TIME OF ONE HOUR AFTER BURST. EXTRA-RELATED BY MEANS OF 1.44 DECAV LAW. THESE VALUES DID NOT ACTUALLY EXIST SINCE TIMES OF ARRIVAL OF CONTAMINANT WERE CONSIDERABLY LATER

170°E

10°E

10°E

BIKAR RC  
BIKAR

12°N

50 40 30 20 10 10°N

50 40 30 20 10 11°N

50 40 30 20 10 12°N

50 40 30 20 10 13°N

50 40 30 20 10 14°N

50 40 30 20 10 15°N

50 40 30 20 10 16°N

50 40 30 20 10 17°N

50 40 30 20 10 18°N

50 40 30 20 10 19°N

50 40 30 20 10 20°N

50 40 30 20 10 21°N

50 40 30 20 10 22°N

50 40 30 20 10 23°N

50 40 30 20 10 24°N

50 40 30 20 10 25°N

50 40 30 20 10 26°N

50 40 30 20 10 27°N

50 40 30 20 10 28°N

50 40 30 20 10 29°N

50 40 30 20 10 30°N

50 40 30 20 10 31°N

50 40 30 20 10 32°N

50 40 30 20 10 33°N

50 40 30 20 10 34°N

50 40 30 20 10 35°N

50 40 30 20 10 36°N

50 40 30 20 10 37°N

50 40 30 20 10 38°N

50 40 30 20 10 39°N

50 40 30 20 10 40°N

CHARLIE FOX  
TARE MAN  
BIKINI

TO BEKORIKALAM  
RONGELAP  
AILINGINAE  
WOTHO

1100 MAIN  
1200 TERN  
1300 TERN  
1400 TERN  
1500 TERN  
1600 TERN  
1700 TERN  
1800 TERN  
1900 TERN  
2000 TERN  
2100 TERN  
2200 TERN  
2300 TERN  
2400 TERN  
2500 TERN  
2600 TERN  
2700 TERN  
2800 TERN  
2900 TERN  
3000 TERN  
3100 TERN  
3200 TERN  
3300 TERN  
3400 TERN  
3500 TERN  
3600 TERN  
3700 TERN  
3800 TERN  
3900 TERN  
4000 TERN  
4100 TERN  
4200 TERN  
4300 TERN  
4400 TERN  
4500 TERN  
4600 TERN  
4700 TERN  
4800 TERN  
4900 TERN  
5000 TERN  
5100 TERN  
5200 TERN  
5300 TERN  
5400 TERN  
5500 TERN  
5600 TERN  
5700 TERN  
5800 TERN  
5900 TERN  
6000 TERN  
6100 TERN  
6200 TERN  
6300 TERN  
6400 TERN  
6500 TERN  
6600 TERN  
6700 TERN  
6800 TERN  
6900 TERN  
7000 TERN  
7100 TERN  
7200 TERN  
7300 TERN  
7400 TERN  
7500 TERN  
7600 TERN  
7700 TERN  
7800 TERN  
7900 TERN  
8000 TERN  
8100 TERN  
8200 TERN  
8300 TERN  
8400 TERN  
8500 TERN  
8600 TERN  
8700 TERN  
8800 TERN  
8900 TERN  
9000 TERN  
9100 TERN  
9200 TERN  
9300 TERN  
9400 TERN  
9500 TERN  
9600 TERN  
9700 TERN  
9800 TERN  
9900 TERN  
10000 TERN

UTIRIK  
TAKA

AILUK

JEMO

LIKIEP

WOTHO

10°N



FIG. 5



Dr. White

Sample Designator:  
 S - SOIL  
 W - WATER  
 F - FROGS  
 U - URINE

Dr. White  
 3-9-83

<u>SAMPLE NO.</u>	<u>DATE</u>	<u>TIME</u>	<u>LOCATION</u>	<u>DESCRIPTION</u>	<u>ANALYST</u>
31	6 March	0800	Makap Island, Makap Atoll	Upper layer bare soil in random spots unsheltered by trees or shrubs, etc.	"
W1	"	"	"	Collected from largest clatarn on heaviest populated island of atoll	"
S2	6 March	1200	Jemo Island	Same as S1	"
W2	"	"	"	Same as W1	"
S3	6 March	1700	Aluk Island, Aluk Atoll	Same as S1	"
W3	"	"	"	Same as W1	"
S4	7 March	1300	Mejit Island	Same as S1	"
W4	"	"	"	Same as W1	"
W5-8	4 March	0900	Utirik Atoll	Composites of 4 water samples taken by the USS REMBELLAN	"
S-5	5 March	1600H	Ormed Island, Kotje Atoll	Composites of 5 samples (1 beach, 3 mid-village, 1 back village)	Dr. T. White
W11	5 March	"	"	Composites: 1/2 from wall, 1/2 from catch-basin	"
S6	5 March	1730	Erkub Island, Kerub Atoll	Composite of 2 samples (1 mid-village, 1 half-way to beach)	"
S7	6 March	1130	Kaven Island, Maloelap Atoll	Composites of 4 samples (2 from village, 2 from paths to beach)	"
W12	"	"	"	1 from wall	"
W13	"	"	"	1 from catch-basin	"
S8	6 March	1630	Motho Island, Motho Atoll	Composites of 3 samples (1 by wall, 2 mid-village)	"

<u>SAMPLE NO.</u>	<u>DATE</u>	<u>TIME</u>	<u>LOCATION</u>	<u>DESCRIPTION</u>	<u>AGENT</u>
19	6 March	1630	Hotho Island, Motu Atooll	1 from well (catch-basin dry for 1 month plus)	Dr. T. Kato
S9	7 March	1200	Dalap Island, Majuro Atooll	Composite of 4 samples (near Admin Bldg)	"
120	"	"	"	Tap water	"
2 S10	"	"	Utirik	Composite of 3 samples	"
121	3 March	0930	Rongelap Island	Composite of 6 bottles Quart included to show location of bottles on Rongelap Island	Dr. P. Scoville
122	8	8	"	Central cistern of village	"
123	8	8	"	Cistern water from north part of Island	"
124	8	8	"	Cistern water from northern most village	"
125	8	8	"	South cistern in village	"
126	9	9	Utirik	Cistern	"
127	9	9	"	Cistern	"
128	10	10	Eniwetak Island	Distillation water Rongelap Atooll	"
121	8	8	Rongelap Island	Soil from north part of Island	"
-S12	8	8	"	Center portion of Island	"
S13	8	8	"	1 mile north of Rongelap village	"
S14	8	8	"	Near south cistern of village	"
S15	8	8	Erllitipu Island	"	"
S16	8	8	Eniwetak Island	"	"
S17	8	8	Kabolio Island	"	"
S18	8	8	Utirik Island	"	"
S19	9	9	Makar Island	"	"

<u>SAMPLE NO.</u>	<u>DATE</u>	<u>TIME</u>	<u>LOCATION</u>	<u>DESCRIPTION</u>
820	10 Mar,		Endvatak Island	
821	10		Sifo Island	Temporary Village - Allingfinae Atoll
822	9		Bikar Island	Rollage, windward side

ACRBT  
 Dr. P. Scoville